



Tyler Thorsen

Norman Loeb, Seiji Kato, Fred Rose

Closure and attribution of the observed variability in the Earth's energy budget

Introduction

- CERES can provide observations of the total TOA energy budget, but isolating individual contributions must rely on bottom-up calculations based on cloud, surface, and atmospheric inputs.
 - Such a decomposition can provide valuable insights into the into the underlying causes of observed changes in the radiation budget.
 - Relating the individual radiative responses observed on short time scales to those from long-term climate change is worthwhile but challenging.

Introduction

- CERES can provide observations of the total TOA energy budget, but isolating individual contributions must rely on bottom-up calculations based on cloud, surface, and atmospheric inputs.
 - Such a decomposition can provide valuable insights into the underlying causes of observed changes in the radiation budget.
 - Relating the individual radiative responses observed on short time scales to those from long-term climate change is worthwhile but challenging.
- First, it is prudent to understand how accurately one can attribute the total variability in the energy budget to the underlying individual radiative anomalies.
- **Extend the analyses of Thorsen et al. (2018) to explore how best to decompose and reproduce the observed EBAF-TOA anomalies over the past 2 decades.**

Thorsen et al. 2018

- Applied partial radiative perturbation (PRP) calculations to observations
- The impact of a perturbation Δx on the flux ($\delta F_{\Delta x}$):

$$\delta F_{\Delta x} = F(x + \Delta x, y_1, \dots, y_N) - F(x, y_1, \dots, y_N)$$

- Fluxes (F) computed with gridded month mean inputs x, y_1, \dots, y_N (Fu-Liou)
- Δx is a gridded deseasonalized monthly mean anomaly of a particular input variable
- ** more complicated than this...

Thorsen et al. 2018

- Applied partial radiative perturbation (PRP) calculations to observations
- The impact of a perturbation Δx on the flux ($\delta F_{\Delta x}$):

$$\delta F_{\Delta x} = F(x + \Delta x, y_1, \dots, y_N) - F(x, y_1, \dots, y_N)$$

- Fluxes (F) computed with gridded month mean inputs x, y_1, \dots, y_N (Fu-Liou)
- Δx is a gridded deseasonalized monthly mean anomaly of a particular input variable
- ** more complicated than this...
- Compute $\delta F_{\Delta x}$ for:
 - ① Temperature (skin/surface + atmosphere),
 - ② Water vapor,
 - ③ Surface albedo/emissivity,
 - ④ Aerosols,
 - ⑤ Insolation and trace gases (ozone, carbon dioxide, methane, nitrous oxide, CFC-11, CFC-12, HCFC-22),
 - ⑥ Cloud fraction,
 - ⑦ Cloud optical depth,
 - ⑧ Other cloud properties (top/base pressure, phase, effective particle size)

Thorsen et al. 2018

- Applied partial radiative perturbation (PRP) calculations to observations
- The impact of a perturbation Δx on the flux ($\delta F_{\Delta x}$):

$$\delta F_{\Delta x} = F(x + \Delta x, y_1, \dots, y_N) - F(x, y_1, \dots, y_N)$$

- Fluxes (F) computed with gridded month mean inputs x, y_1, \dots, y_N (Fu-Liou)
- Δx is a gridded deseasonalized monthly mean anomaly of a particular input variable
- ** more complicated than this...
- Compute $\delta F_{\Delta x}$ for:
 - ① Temperature (skin/surface + atmosphere), ② Water vapor,
 - ③ Surface albedo/emissivity, ④ Aerosols,
 - ⑤ Insolation and trace gases (ozone, carbon dioxide, methane, nitrous oxide, CFC-11, CFC-12, HCFC-22),
 - ⑥ Cloud fraction, ⑦ Cloud optical depth,
 - ⑧ Other cloud properties (top/base pressure, phase, effective particle size)
- Sum ⑥+⑦+⑧ = total impact of cloud anomalies $\delta F_{\Delta cld}^{PRP}$: “pure PRP approach” for isolating cloud radiative impact

Adjusted CRE approach

- Alternative to the pure PRP approach, the impact of clouds can be isolated by adjusting the cloud radiative effect (CRE, the difference in the all-sky and clear-sky radiative flux) for the masking effects of non-cloud variables (Soden et al. 2008):

$$\delta F_{\Delta cld}^{CRE} = \Delta CRE^{EBAF} - \sum_i [\delta F_{x_i}(\text{all-sky}) - \delta F_{x_i}(\text{clear-sky})]$$

- ΔCRE^{EBAF} : monthly gridded deseasonalized CRE anomalies from EBAF-TOA
- $\delta F_{x_i}(\text{all-sky})$ and $\delta F_{x_i}(\text{clear-sky})$ are the non-cloud PRP calculations

Input datasets

- 1 degree gridded monthly mean data from March 2000 through March 2020
- Non-cloud properties: GEOS-5.4.1 (temperature, water vapor, ozone), MATCH/MODIS (aerosols), AIRS L3 (CO₂, CH₄), NOAA ESRL (N₂O, CFCs), CERES SAH (surface albedo), SORCE (insolation)

Input datasets

- 1 degree gridded monthly mean data from March 2000 through March 2020
- Non-cloud properties: GEOS-5.4.1 (temperature, water vapor, ozone), MATCH/MODIS (aerosols), AIRS L3 (CO₂, CH₄), NOAA ESRL (N₂O, CFCs), CERES SAH (surface albedo), SORCE (insolation)
- Clouds:
 - ❶ SYN (Ed4): MODIS+GEO diurnally-resolved properties [used in Thorsen et al. 2018]
 - ❷ Terra and Aqua SSF (Ed4): no (potential) geostationary artifacts (MODIS-only)

Input datasets

- 1 degree gridded monthly mean data from March 2000 through March 2020
- Non-cloud properties: GEOS-5.4.1 (temperature, water vapor, ozone), MATCH/MODIS (aerosols), AIRS L3 (CO₂, CH₄), NOAA ESRL (N₂O, CFCs), CERES SAH (surface albedo), SORCE (insolation)
- Clouds:
 - ❶ SYN (Ed4): MODIS+GEO diurnally-resolved properties [used in Thorsen et al. 2018]
 - ❷ Terra and Aqua SSF (Ed4): no (potential) geostationary artifacts (MODIS-only)
- Also explore using a “tuned” set of inputs from the EBAF-surface algorithm
 - Produces 2 optimized sets of cloud, surface, and atmospheric inputs: (1) those tuned to all-sky EBAF-TOA fluxes, (2) those tuned to clear-sky EBAF-TOA fluxes

Input datasets

- 1 degree gridded monthly mean data from March 2000 through March 2020
- Non-cloud properties: GEOS-5.4.1 (temperature, water vapor, ozone), MATCH/MODIS (aerosols), AIRS L3 (CO₂, CH₄), NOAA ESRL (N₂O, CFCs), CERES SAH (surface albedo), SORCE (insolation)
- Clouds:
 - ❶ SYN (Ed4): MODIS+GEO diurnally-resolved properties [used in Thorsen et al. 2018]
 - ❷ Terra and Aqua SSF (Ed4): no (potential) geostationary artifacts (MODIS-only)
- Also explore using a “tuned” set of inputs from the EBAF-surface algorithm
 - Produces 2 optimized sets of cloud, surface, and atmospheric inputs: (1) those tuned to all-sky EBAF-TOA fluxes, (2) those tuned to clear-sky EBAF-TOA fluxes
- Using these tuned inputs in our PRP calculations will presumably provide a more accurate decomposition of the EBAF-TOA variability
 - Highly under-constrained problem: possible to not find the optimal solution, uncorrected biases
→ insight from PRP calculations.

Comparisons to EBAF-TOA

- Comparisons to EBAF Ed4.1 TOA data: absorbed solar radiation (ASR), longwave (LW) radiation, and net (ASR+LW). TOA fluxes are defined positive downwards (LW = -OLR)
- EBAF-TOA anomaly time series compared to bottom-up PRP calculations of the total radiative anomalies, i.e the sum of all the individual (x_i) PRP calculations:

$$\delta F_{\Delta\text{total}} = \sum_i \delta F_{\Delta x_i}.$$

Comparisons to EBAF-TOA

- Comparisons to EBAF Ed4.1 TOA data: absorbed solar radiation (ASR), longwave (LW) radiation, and net (ASR+LW). TOA fluxes are defined positive downwards (LW = -OLR)
- EBAF-TOA anomaly time series compared to bottom-up PRP calculations of the total radiative anomalies, i.e the sum of all the individual (x_i) PRP calculations:

$$\delta F_{\Delta\text{total}} = \sum_i \delta F_{\Delta x_i}.$$

- Compute metrics for (1) global mean time series, (2) regional [1x1 degree] time series
 - ① Coefficient of determination (R^2)
 - ② Residual trend: i.e. trend in $\delta F_{\Delta\text{total}} - \Delta F^{EBAF}$

Comparisons to EBAF-TOA

- Comparisons to EBAF Ed4.1 TOA data: absorbed solar radiation (ASR), longwave (LW) radiation, and net (ASR+LW). TOA fluxes are defined positive downwards (LW = -OLR)
- EBAF-TOA anomaly time series compared to bottom-up PRP calculations of the total radiative anomalies, i.e the sum of all the individual (x_i) PRP calculations:


$$\delta F_{\Delta\text{total}} = \sum_i \delta F_{\Delta x_i}.$$

- Compute metrics for (1) global mean time series, (2) regional [1x1 degree] time series
 - ① Coefficient of determination (R^2)
 - ② Residual trend: i.e. trend in $\delta F_{\Delta\text{total}} - \Delta F^{EBAF}$
- Make comparisons for PRP calculations based on the various combinations of:
 - ① Cloud inputs: SYN, SSF
 - ② EBAF-surface tuned inputs: untuned (0), all-sky tuned (AT), clear-sky tuned (CT)
 - ③ Approach for isolating clouds anomalies: pure PRP, adjusted CRE

Variability closure (global mean time series)

Pure PRP ($\delta F_{\Delta cld}^{PRP}$):

- SYN^0 : low R^2 in the LW, better for ASR




Variability closure

(global mean time series)

Pure PRP ($\delta F_{\Delta cld}^{PRP}$):

- SYN^0 : low R^2 in the LW, better for ASR
- SYN^{AT} : large improvements from all-sky tuned inputs (cuts the residual trends in about half, large R^2 values)




Variability closure

(global mean time series)


Pure PRP ($\delta F_{\Delta cld}^{PRP}$):

- SYN^0 : low R^2 in the LW, better for ASR
- SYN^{AT} : large improvements from all-sky tuned inputs (cuts the residual trends in about half, large R^2 values)
- SSF^0 : mostly better comparisons than SYN^0 , but not better than SYN^{AT}




Impact of all-sky tuning (i.e. insight into EBAF-surface algorithm)

- (a) \rightarrow (b): geostationary artifacts greatly reduced in the residual trends (EBAF-surface algorithm is improving agreement to EBAF-TOA)




Impact of all-sky tuning (i.e. insight into EBAF-surface algorithm)

- (a) \rightarrow (b): geostationary artifacts greatly reduced in the residual trends (EBAF-surface algorithm is improving agreement to EBAF-TOA)
- This reduction in the residual trend is mostly achieved by adjusting the cloud properties and water vapor



Impact of all-sky tuning (i.e. insight into EBAF-surface algorithm)

- (a) \rightarrow (b): geostationary artifacts greatly reduced in the residual trends (EBAF-surface algorithm is improving agreement to EBAF-TOA)
- This reduction in the residual trend is mostly achieved by adjusting the cloud properties and water vapor
- However, the water vapor radiative trends in SYN^{AT} (f) have geostationary artifacts
- EBAF-surface algorithm is erroneously adjusting the water vapor to compensate for uncertainties in the cloud properties \rightarrow best to avoid using all-sky tuned inputs




Variability closure

(global mean time series)

Pure PRP ($\delta F_{\Delta cld}^{PRP}$):

- Avoid **SYN^{AT}**, but can still leverage clear-sky tuned inputs




Variability closure

(global mean time series)

Pure PRP ($\delta F_{\Delta cld}^{PRP}$):

- Avoid SYN^{AT} , but can still leverage clear-sky tuned inputs
- SYN^{CT} / SSF^{CT} : not always an improvement over SYN^0 / $SSF^0 \rightarrow$ revealing of cloud errors that were radiatively masked in the untuned inputs




Variability closure

(global mean time series)

Pure PRP ($\delta F_{\Delta cld}^{PRP}$):

- Avoid SYN^{AT} , but can still leverage clear-sky tuned inputs
- SYN^{CT} / SSF^{CT} : not always an improvement over SYN^0 / $SSF^0 \rightarrow$ revealing of cloud errors that were radiatively masked in the untuned inputs

Adjusted CRE ($\delta F_{\Delta cld}^{CRE}$):




Variability closure

(global mean time series)

Pure PRP ($\delta F_{\Delta cld}^{PRP}$):

- Avoid SYN^{AT} , but can still leverage clear-sky tuned inputs
- SYN^{CT} / SSF^{CT} : not always an improvement over SYN^0 / $SSF^0 \rightarrow$ revealing of cloud errors that were radiatively masked in the untuned inputs

Adjusted CRE ($\delta F_{\Delta cld}^{CRE}$):




Variability closure

(global mean time series)

Pure PRP ($\delta F_{\Delta cld}^{PRP}$):

- Avoid SYN^{AT} , but can still leverage clear-sky tuned inputs
- SYN^{CT} / SSF^{CT} : not always an improvement over SYN^0 / $SSF^0 \rightarrow$ revealing of cloud errors that were radiatively masked in the untuned inputs

Adjusted CRE ($\delta F_{\Delta cld}^{CRE}$):



Variability closure


(global mean time series)

Pure PRP ($\delta F_{\Delta cld}^{PRP}$):

- Avoid SYN^{AT} , but can still leverage clear-sky tuned inputs
- SYN^{CT} / SSF^{CT} : not always an improvement over SYN^0 / $SSF^0 \rightarrow$ revealing of cloud errors that were radiatively masked in the untuned inputs

Adjusted CRE ($\delta F_{\Delta cld}^{CRE}$):

- In general, the adjusted CRE approach (solid box) performs better than the pure PRP approach (dashed box), particularly when the clear-sky tuned inputs are used



Variability closure


(global mean time series)

Pure PRP ($\delta F_{\Delta cld}^{PRP}$):

- Avoid SYN^{AT} , but can still leverage clear-sky tuned inputs
- SYN^{CT} / SSF^{CT} : not always an improvement over SYN^0 / $SSF^0 \rightarrow$ revealing of cloud errors that were radiatively masked in the untuned inputs

Adjusted CRE ($\delta F_{\Delta cld}^{CRE}$):

- In general, the adjusted CRE approach (solid box) performs better than the pure PRP approach (dashed box), particularly when the clear-sky tuned inputs are used
- Similar comparisons regardless of the cloud properties used + the individual radiative anomalies are very similar as well (not shown) \rightarrow **adjusted CRE approach is not sensitive to the choice of cloud properties**



Variability closure


(global mean time series)

Pure PRP ($\delta F_{\Delta cld}^{PRP}$):

- Avoid SYN^{AT} , but can still leverage clear-sky tuned inputs
- SYN^{CT} / SSF^{CT} : not always an improvement over SYN^0 / $SSF^0 \rightarrow$ revealing of cloud errors that were radiatively masked in the untuned inputs

Adjusted CRE ($\delta F_{\Delta cld}^{CRE}$):

- In general, the adjusted CRE approach (solid box) performs better than the pure PRP approach (dashed box), particularly when the clear-sky tuned inputs are used
- Similar comparisons regardless of the cloud properties used + the individual radiative anomalies are very similar as well (not shown) \rightarrow **adjusted CRE approach is not sensitive to the choice of cloud properties**
- The combination of the **adjusted CRE approach and clear-sky tuned inputs** is the only set of calculations that gives **statistically insignificant residual trends** in the ASR, LW, and net flux anomalies



Conclusions: variability closure

Conclusions: variability closure

- All-sky tuned inputs (from the EBAF-surface algorithm) improve agreement to EBAF-TOA, but this is partly achieved by adjusting water vapor to compensate for uncertainties the SYN cloud properties.

Conclusions: variability closure

- All-sky tuned inputs (from the EBAF-surface algorithm) improve agreement to EBAF-TOA, but this is partly achieved by adjusting water vapor to compensate for uncertainties the SYN cloud properties.
- Isolating cloud anomalies using the adjusted CRE approach performs better than the pure PRP calculations, especially when paired with clear-sky tuned inputs.

Conclusions: variability closure

- All-sky tuned inputs (from the EBAF-surface algorithm) improve agreement to EBAF-TOA, but this is partly achieved by adjusting water vapor to compensate for uncertainties the SYN cloud properties.
- Isolating cloud anomalies using the adjusted CRE approach performs better than the pure PRP calculations, especially when paired with clear-sky tuned inputs.
- The adjusted CRE approach:
 - Provides a more direct link to the observed radiative effect of clouds, avoids the need to directly forward model the impact of clouds properties
 - Very insensitive to the cloud properties used (only uses them in a relative sense) → bodes well for assessments based on GCM radiative kernels

Conclusions: variability closure

- All-sky tuned inputs (from the EBAF-surface algorithm) improve agreement to EBAF-TOA, but this is partly achieved by adjusting water vapor to compensate for uncertainties the SYN cloud properties.
- Isolating cloud anomalies using the adjusted CRE approach performs better than the pure PRP calculations, especially when paired with clear-sky tuned inputs.
- The adjusted CRE approach:
 - Provides a more direct link to the observed radiative effect of clouds, avoids the need to directly forward model the impact of clouds properties
 - Very insensitive to the cloud properties used (only uses them in a relative sense) → bodes well for assessments based on GCM radiative kernels
 - But, cannot break down the cloud anomaly further (e.g. cloud property/type) like the PRP approach can → limited insight into the individual cloud property radiative anomalies

Conclusions: variability closure


- All-sky tuned inputs (from the EBAF-surface algorithm) improve agreement to EBAF-TOA, but this is partly achieved by adjusting water vapor to compensate for uncertainties the SYN cloud properties.
- Isolating cloud anomalies using the adjusted CRE approach performs better than the pure PRP calculations, especially when paired with clear-sky tuned inputs.
- The adjusted CRE approach:
 - Provides a more direct link to the observed radiative effect of clouds, avoids the need to directly forward model the impact of clouds properties
 - Very insensitive to the cloud properties used (only uses them in a relative sense) → bodes well for assessments based on GCM radiative kernels
 - But, cannot break down the cloud anomaly further (e.g. cloud property/type) like the PRP approach can → limited insight into the individual cloud property radiative anomalies
 - Computed CRE is forced to be the same as the EBAF-TOA CRE → best to use the cloud properties that gives a computed CRE most similar to EBAF-TOA CRE (SSF clouds)

Conclusions: variability closure

- All-sky tuned inputs (from the EBAF-surface algorithm) improve agreement to EBAF-TOA, but this is partly achieved by adjusting water vapor to compensate for uncertainties the SYN cloud properties.
- Isolating cloud anomalies using the adjusted CRE approach performs better than the pure PRP calculations, especially when paired with clear-sky tuned inputs.
- The adjusted CRE approach:
 - Provides a more direct link to the observed radiative effect of clouds, avoids the need to directly forward model the impact of clouds properties
 - Very insensitive to the cloud properties used (only uses them in a relative sense) → bodes well for assessments based on GCM radiative kernels
 - But, cannot break down the cloud anomaly further (e.g. cloud property/type) like the PRP approach can → limited insight into the individual cloud property radiative anomalies
 - Computed CRE is forced to be the same as the EBAF-TOA CRE → best to use the cloud properties that gives a computed CRE most similar to EBAF-TOA CRE (SSF clouds)


**To decompose the energy budget, it is best to use calculations based on
SSF cloud properties, clear-sky tuned inputs,
and the adjusted CRE approach to isolate cloud anomalies.**

Attribution




- More in Norman's talk tomorrow...

Attribution




- More in Norman's talk tomorrow...
- (a) Positive ASR trend in EBAF-TOA is mostly the result of cloud and surface albedo changes

Attribution



- More in Norman's talk tomorrow...
- (a) Positive ASR trend in EBAF-TOA is mostly the result of cloud and surface albedo changes
- (b) The negative EBAF-TOA LW trend is due to increased OLR from increasing temperatures and cloud changes which is partly offset by contributions from water vapor and trace gases

Attribution



- More in Norman's talk tomorrow...
- (a) Positive ASR trend in EBAF-TOA is mostly the result of cloud and surface albedo changes
- (b) The negative EBAF-TOA LW trend is due to increased OLR from increasing temperatures and cloud changes which is partly offset by contributions from water vapor and trace gases
- (c) Positive net trend with the underlying changes all contributing significantly with the exception of aerosols. Negative contribution from temperature is overwhelmed by the positive contributions from clouds, water vapor, surface albedo, and trace gas changes.